

MYCOLOGICAL SURVEY OF RÍO CAMUY CAVES PARK, PUERTO RICO

ÁNGEL M. NIEVES-RIVERA

Department of Marine Sciences, P.O. Box 9013, University of Puerto Rico, Mayagüez PR 00681-9013 USA anieves@coqui.net

The Río Camuy Caves Park is part of the Río Camuy Cave System in the northern karst belt of Puerto Rico. Thirty-nine fungal species and 11 cellular and plasmodial slime molds are new records for caves and sinkholes of the Río Camuy Cave Park. Two of the cellular slime molds Dictyostelium citrinum and D. macrocephalum are new records for Puerto Rico.

El Parque de las Cavernas del Río Camuy pertenece al Sistema de Cavernas del Río Camuy en el cinturón del carso norteño de Puerto Rico. Se informan 39 especies de hongos y 11 hongos mucilaginosos celulares y plasmodiales como nuevos registros para cuevas y sumideros del Parque de las Cavernas del Río Camuy. Dos de los hongos mucilaginosos celulares Dictyostelium citrinum y D. macrocephalum son nuevos registros para Puerto Rico.

Fungi play roles in cave communities as saprotrophs or pathogens. Fungi affect the population dynamics of cave biota and contribute in the decomposition of organic matter, making it available to other members of the cave community. Diets of many insect, including cavernicoles, include fungal spores and mycelia (Culver 1982), and spores from at least 15 fungal genera probably constitute an important element in the diet of springtails (Collembola: Insecta) (Cubbon 1976; Culver 1982; Roselló *et al.*, 1986). Insects are considered an effective dispersion vector of fungal spores and other microorganisms because of nonspecific factors in attachment to host insect cuticles (Boucias *et al.* 1988). Chiropterans, insects, and rodents transport many fungal species to this bat-guano enriched soil environments (Hoff & Bigler 1981).

The Río Camuy Cave System [RCCS] is a 15-km long, subtropical karst feature carved by the Camuy River in the subtropical moist forest (Ewel & Whitmore 1973) of northern Puerto Rico. This system, one of the largest caves in Puerto Rico, is part of the National Parks Company of the Commonwealth of Puerto Rico natural protected area. In the 1960s, Russell and Jeanne Gurnee described the limestone features, caves, conservation, and future development, of what is now Río Camuy Caves Park [RCCP] (Gurnee 1967; Gurnee & Gurnee 1974, 1987; Martínez-Oquendo 1983) (Fig. 1). The geology and hydrology of the RCCS were treated by Thraillkill (1967), Monroe (1976, 1980), Torres-González (1983), Troester and White (1986), and Troester (1988, 1994). A recent study (Lugo *et al.* 2001) summarized what is known about the karst of Puerto Rico (including RCCS), including its importance and anthropogenic impact. Bat-guano enriched soil was formerly mined from Puerto Rican caves (Cadilla 1962; Beck *et al.* 1976; Campbell 1994; Frank 1998), but the guano in Río Camuy caves apparently was not heavily exploited (Cardona-Bonet, pers. comm. 2001).

In general, the rainy season in Puerto Rico peaks in May-June and August-September. Air temperatures in RCCS range from 18-25° C (Mercado, pers. comm. 2000), relative humidity from 88-100% (Troester 1988), and the average annual rain-




Figure 1. Map of Río Camuy Caves Park. Redrawn after Luis Ayala (November 1988) by permission (National Parks Company of the Commonwealth of Puerto Rico), based on Gurnee & Gurnee (1974) original map.

fall for Lares, 7 km SW of RCCP, is 2435 mm.

The vascular flora of the sinkholes and mogotes of RCCP was surveyed by Del Llano (1982) and Vives *et al.* (1985), recording 136 and 73 plant species, respectively. Reyes-Colón (1999) and Reyes-Colón and Sastre-DJ (2000) recorded 50

species: 31 mosses (11% of the moss flora of Puerto Rico) and 19 liverworts (8% of the liverwort flora of Puerto Rico) from 2 sinkholes (Espiral and Empalme Sinks) in RCCP.

The caves and sinkholes of RCCP are inhabited by a variety of invertebrates and vertebrates (Sullivan 1966, 1967; Peck 1974, 1981, 1994; Díaz-Díaz 1982, 1983; Santos-Flores 2001; Lugo *et al.* 2001). Multispecies assemblages of bats are reported from RCCP, as is common in many Puerto Rican caves (Rodríguez-Durán & Lewis 1987; Conde Costas & González 1990; Rodríguez-Durán 1998). With the exception of the endemic amphipod *Allovecelia gurneei* Holsinger & Peck (Holsinger & Peck 1967) and the cave cockroach *Nelipophygus* sp. (Peck 1981), no other troglobites have been reported from RCCS.

Betancourt *et al.* (1988) recorded 8 genera of mitosporic fungi (*Aspergillus* sp., *Cladosporium* sp., *Curvularia* sp., *Fusarium* sp., *Geotrichum* sp., *Scopulariopsis* sp., *Sepedonium* sp., and *Rhizopus* sp.) from Ensueño Cave (3 km south of RCCP). Eight species of aquatic hyphomycetes (*Brachyosphaera jamaicensis* [Crane & Dumont] Descals, *Clavariopsis azlanii* Nawawi, *Dendrosporium lobatum* Plakidas & Edgerton: Crane, *Isthmolongispora quadricellularia* Matsushima, *Tricladiospora brunnea* [Nawawi] Nawawi & Kuth., *Tripodermum* sp., *Varicosporium giganteum* Crane, *Xenosporium berkeleyi* [Curtis] Pirozynski) were also isolated in river foam from Tres Pueblos Sinkhole (Santos-Flores & Betancourt-López 1997) in RCCP.

This report summarizes a mycological survey of the bat-guano enriched soil, leaf litter, wood, top soil, streams, and intermittent pools in RCCP.

METHODS

I conducted a basic inventory of the fungi, and the cellular/plasmoidal slime molds at Cathedral Cave, Clara Cave, Empalme Sinkhole, and Tres Pueblos Sinkhole (RCCP, Bayaney topographic quadrangle) (Figure 1). These sites are described below.

•Cathedral Cave (“Cueva La Catedral”) (18°21’04.2” N, 66°48’99.3” W), a hydrological inactive cave, is located adjacent to the hydrographic basin of the Río Camuy, at the north end of RCCP (Fig. 1). This cave is 122 m long, with 2 main parallel galleries (or chambers), which run north-south and are connected by a short passage, with 2 natural skylights, about 24 and 38 m above the floor of the cave. The west hall contains rock art of presumed aboriginal origin (42 pictographs in total; Nieves-Rivera, unpubl. data).

•Clara Cave [“Cueva Clara de Empalme”, “Empalme Cave” of Peck (1974)] (18°20’85.4” N, 66°49’16.0” W) is located south of Cathedral Cave (Fig. 1). This cave is part of the ancient course of the Río Camuy and is higher than the present river level. The main passage is a diagonal tunnel 115 m long and 52 m high. A secondary passage extends 30 m to the west. The primary entrance is shown in Figure 2A; there is also

a vertical shaft entrance at Empalme Sinkhole. Previous hydrological studies of 2 sites (streamlet/pool R-1 [Fig. 2B] and pool R-2, both intermittent) of Cueva Clara reported concentrations of 7.9-8.6 mg/L of dissolved oxygen at 22°C, and 0.131 to 0.040 mg/L of total phosphorous, respectively (Díaz-Díaz 1982). Pool water pH has been measured between 9.1 and 9.2 with conductivity rendering 200 μ (Nieves-Rivera, unpubl. data). Fecal coliform counts have been reported as 1160 cells/mL for R-1 and 1810 cells/mL for R-2, respectively, indicating a runoff water with high organic input (fecal matter and detritus) (Díaz-Díaz 1982).

•Empalme Sinkhole (“Sumidero de Empalme”) (18°20’79.9” N, 66°49’11.2” W) is a 99-m wide doline with a 125-m deep vertical shaft, connecting directly to Clara Cave (Fig. 1) and to the conical-shaped “Cueva Alta del Norte” (Fig. 2C). At the bottom, and along the upper reaches of the Río Camuy, numerous collapsed blocks are present, along with top soil composed of clay or dirt, leaf litter, mosses, and rocks with lichens. Further description of this doline was given by Gurnee (1967), Gurnee & Gurnee (1974), Reyes-Colón (1999), and Reyes-Colón & Sastre-DJ (2000).


•Tres Pueblos Sinkhole (“Sumidero Tres Pueblos”) (18°20’30.5” N, 66°49’26.3” W) is a doline in RCCP, and was described by Gurnee (1967) and Gurnee & Gurnee (1974).

FIELD WORK


Field work at Cathedral Cave, Clara Cave, and Empalme Sinkhole was carried out on 8 August 1999, supplemented by river water and mud samples collected from Tres Pueblos Sinkhole in May 1993. Undergraduate and graduate students, and professors from the University of Puerto Rico at Mayagüez Campus (Departments of Crop Protection and Marine Sciences) participated in the survey.

LABORATORY WORK

Fungi were isolated from field-collected samples. Samples of bat-guano enriched soil, leaf litter, wood, water, and soil were collected using sterile plastic bags (Whirl Pak). Higher fungi (ascomycetes and basidiomycetes) were collected and processed following Nieves-Rivera *et al.* (1999). Bat-guano enriched soil samples were processed according to Orpurt (1964), Gaur and Lichtwardt (1980), and Rutherford and Huang (1994). Freshwater fungi sampling followed Sparrow (1960), Nieves-Rivera and Betancourt-López (1994) and Santos-Flores and Betancourt-López (1997). Potato dextrose agar acidified with 10% lactic acid, Rose Bengal agar, and Sabouraud’s dextrose agar were used as fungal growth media. Wet mount observations were made with a Nikon Labophoto-2 microscope. All voucher specimens were taken to the Center for Forest Mycology (U.S. Department of Agriculture, Forest Products Laboratory, Sabana Station, Puerto Rico) where they are being prepared for deposition in the herbarium of the Department of Natural Sciences, University of Puerto Rico



Piedras, Puerto Rico.



Figures 2A-C. Río Camuy Caves Park study sites. A. Main entrance of Clara Cave. B. Intermittent streamlet R-1 seen from inside Clara Cave. C. Inside Empalme Sinkhole (with a view of Cueva Alta del Norte), mesophytic forest, Río Camuy is at the bottom.

RESULTS AND DISCUSSION

A total of 45 true-fungi (39 new records) and 16 fungal-like organisms (11 Dictyosteliomycota/Myxomycota and 5 Oomycota; all new records) are reported from RCCS. Of these, a total of 36 taxa (34 micromycetes and 2 macromycetes) from 17 genera were isolated from cave bat-guano enriched soil or

topsoil (Table 1). *Aspergillus*, *Penicillium*, *Fusarium*, *Cladosporium*, *Trichoderma*, *Paecilomyces*, *Curvularia*, *Gliocladium*, *Hirsutella*, and *Sepedonium* were the most common hyphomycete genera in cave soil (Table 1). All these genera are dematiaceous hyphomycetes. *Mycelia Sterilia sensu* Rutherford & Huang (1994) was also present in cave soil. Mitosporic fungi isolated in this study were present as spores, actively growing mycelium or as resting mycelial cysts. Other frequently isolated fungi were members of the Zygomycota, *Mucor*, *Rhizopus*, and an undetermined entomophylic species. The pathogen *Histoplasma capsulatum* Darling, the causative agent of histoplasmosis, was not isolated from cave samples, but was found in 1979 in Humo Cave, part of RCCS (Carvajal-Zamora, pers. comm. 1998; Table 1).

The habitat below the skylights and at other openings of RCCS differs little from the surface (Gillieson 1996). Outside of the study area, *Xylaria* (Ascomycota) and *Lepiota* (Basidiomycota) have been found in caves below skylights in Espinar Cave (Mona Island), De los Santos/Braceros Cave

Table 1. Summary of the mycobiota of the Río Camuy Cave System, Puerto Rico. Nomenclature is based on Hawksworth *et al.* (1995), Alexopoulos *et al.* (1996), and Lodge (1996).

TAXON FUNGI [= MYCOTA] HYPHOMYCETES [= MITOSPORIC FUNGI]	FAMILY	SUBSTRATE ¹	REFERENCES ²
<i>Aspergillus flavus</i> Link	Anamorphic (A.) Trichocomaceae	S	2, 5, 7
<i>Aspergillus fumigatus</i> Fresenius	A. Trichocomaceae	S	2, 5, 7
<i>Aspergillus nidulans</i> (Eidam) G. Winter	A. Trichocomaceae	S	2, 5
<i>Aspergillus niger</i> var. <i>niger</i> van Tieghem	A. Trichocomaceae	S	1, 2, 4, 5, 7
<i>Brachiosphaera jamaicensis</i> (Crane & Dumont) Descals	_____	S	6
<i>Cladosporium cladosporioides</i> (Fresenius) deVries	A. Mycosphaerellaceae	D, S, W	2, 5, 7
<i>Cladosporium herbarum</i> (Pers.) Link	A. Mycosphaerellaceae	S, D	1, 2, 5, 7
<i>Clavariopsis azlanii</i> Nawawi	_____	L, R	6
<i>Curvularia</i> sp.	A. Pleosporaceae	S	1, 5
<i>Dendrosporium lobatum</i> Plakidas & Edgerton: Crane	_____	L, R	6
<i>Fusarium</i> cf. <i>oxysporum</i> Schl.	A. Hypocreaceae	S	1, 2, 5, 7
<i>Fusarium solani</i> (Martius) Sacc.	A. Hypocreaceae	S	5, 7
<i>Geotrichum</i> sp.	A. Dipodascaceae	S	1, 5
<i>Gliocladium</i> cf. <i>roseum</i> Bain.	_____	S	2, 5
<i>Hirsutella</i> sp.	_____	I	2, 5
<i>Histoplasma capsulatum</i> Darling ³	A. Onygenaceae	S	3, 7
<i>Isthmolongispora quadricellularia</i> Matsushima	_____	L, R	6
<i>Neurospora crassa</i> Shear & Dodge	Sordariaceae	S	2, 5, 7
<i>Paecilomyces</i> sp.	A. Trichocomaceae	S	2, 5
<i>Penicillium</i> cf. <i>crysogenum</i> Thom	_____	S	2, 5, 7
<i>Penicillium lilacinum</i> Thom	_____	S	2, 5, 7
<i>Penicillium roqueforti</i> Thom	_____	S	2, 5
<i>Penicillium variable</i> Westling	_____	S	2, 5
<i>Scopulariopsis</i> sp.	A. Microascaceae	S	1
<i>Sepedonium</i> sp.	A. Hypocreaceae	S	1, 5
<i>Trichladisopora brunnea</i> (Nawawi) Nawawi & Kuth.	_____	L, R	6
<i>Trichoderma koeningii</i> Oudemans	A. Hypocreaceae	S	2, 5
<i>Trichoderma viridae</i> Pers.: Fr.	A. Hypocreaceae	S	2, 5, 7
<i>Tripodermum</i> sp.	A. Capnodiaceae	L, R	6
<i>Varicosporium giganteum</i> Crane	_____	L, R	6
<i>Xenosporium berkeleyi</i> (Curtis) Pirozynski	_____	L, R	6
<i>Mycelia Sterilia</i>	Agonomycetaceae	S, W, I	2, 5
ASCOMYCOTA			
<i>Xylaria polymorpha</i> (Pers.: Mérat) Grev.	Xylariaceae	W	2, 5, 7
<i>Xylaria</i> sp.	Xylariaceae	W	2, 5
BASIDIOMYCOTA			
<i>Cotylidia aurantica</i> (Pers.) A. L. Welden	Podoscyphaceae	W	8
<i>Geastrum</i> cf. <i>saccatum</i> (Fr.) Ed. Fischer ⁴	Geastraceae	S	5
<i>Lepiota</i> sp.	Agaricaceae	S	2, 5
<i>Marasmius</i> cf. <i>atrorubens</i> (Berk.) Berk.	Tricholomataceae	L	5
<i>Polyporus tenuiculus</i> (Beauv.) Fr. ⁵	Polyporaceae/Coriolaceae	W	5
<i>Ramaria</i> sp.	Ramariaceae	L	8
<i>Rigidoporus microporus</i> (Fr.) Overeem.	Polyporaceae/Coriolaceae	W	8
ZYGOMYCOTA			
<i>Mucor</i> sp.	Mucoraceae	D, S, W	2, 5
<i>Rhizopus stolonifer</i> (Ehr.: Fr.) Vuill.	Mucoraceae	D, S	1, 2, 5, 7
Undetermined species	Mucoraceae	I	5
CHYTRIDIOMYCOTA			
<i>Rhizophydium</i> sp.	Chytridiaceae	P	5
CHROMISTA [= STRAMINIPILA]			
OOMYCOTA			
<i>Achlya americana</i> Humphrey	Saprolegniaceae	R	4, 5, 7
<i>Achlya parasitica</i> Coker	Saprolegniaceae	R	4, 5
<i>Apodachlya</i> sp.	Leptomitaceae	R	4, 5
<i>Brevilegnia</i> sp.	Saprolegniaceae	R	4, 5
<i>Saprolegnia</i> sp.	Saprolegniaceae	R	4, 5

PROTOZOA

DICTYOSTELIOMYCOTA

<i>Dictyostelium aureostipes</i> Cavender, Raper & Norberg ⁶	Dictyosteliaceae	L, S	4, 5
<i>Dictyostelium citrinum</i> Vadell, Holmes & Cavender ⁶	Dictyosteliaceae	L, S	4, 5
<i>Dictyostelium giganteum</i> Singh ⁶	Dictyosteliaceae	L, S	4, 5
<i>Dictyostelium macrocephalum</i> Hagiwara, Yeh & Chien ⁶	Dictyosteliaceae	L, S	4, 5
<i>Dictyostelium mucoroides</i> Brefeld ⁶	Dictyosteliaceae	L, S	4, 5
<i>Dictyostelium purpureum</i> Olive ⁶	Dictyosteliaceae	L, S	4, 5
<i>Polysphondylium pallidum</i> Olive ⁶	Dictyosteliaceae	L, S	4, 5
<i>Polysphondylium violaceum</i> Brefeld ⁶	Dictyosteliaceae	L, S	4, 5

MYXOMYCOTA

<i>Arcyria</i> cf.	Arcyriaceae	B, W	4, 5
<i>Comatricha</i> cf.	Stemonitidaceae	L	5
<i>Stemonitis</i> cf. <i>herbatica</i> Peck	Stemonitidaceae	W	4, 5, 7

¹ Substrates: B = Bat bones; D = Dung; I = Dead insects; L = Leaf litter; P = Pollen; R = River water or foam; S = Bat-guano enriched soil or top soil; W = dead wood. The substrates were the location where the fungus was found growing or just the site from which a spore was isolated.

² References: 1 = Betancourt *et al.* (1988); 2 = Carvajal-Zamora & Nieves-Rivera (1998); 3 = Lewis (1989); 4 = Nieves-Rivera & Carvajal-Zamora (2000); 5 = Nieves-Rivera (present work); 6 = Santos-Flores & Betancourt-López (1997); 7 = Stevenson (1975); 8 = D. J. Lodge (pers. comm. 2001).

³ Carvajal-Zamora (pers. comm. to Norman Veve Carreras [SEPRI] in 1979 and to me in 1998) isolated this pathogen from bat-guano of Humo Cave (= Cueva Humo) in 1979.

⁴ *Geastrum* cf. *saccatum* was collected directly from top soil (not in bat-guano enriched soil), next to the edge of Tres Pueblos Sinkhole.

⁵ *Polyporus tenuiculus* (= *Favolus brasiliensis* [Fr.] Fr.) basidiomata was consumed by *Caracolus caracola* (Pulmonata: Gastropoda) in a similar way as reported by Lodge (1996).

⁶ Species cultured and identified by John C. Landolt, Shepherd College, Shepherdstown, WV.

(Cabo Rojo), and Cathedral Cave (Nieves-Rivera, unpubl. data). These openings admit plant debris from which fungi sporulate and fruiting bodies may develop.

Food is, in general, relatively scarce in deeper portions of caves (Culver 1982; Gillieson 1996). Most of the fungi isolated in this study are edaphic microfungi. These fungi feed and sporulate on different substrates (e.g., a carcasses, hair, seeds, or bat-guano). Use of similar substrata by both lower and higher fungi has been reported for Diablo Cave (Mexico) (Hoffman *et al.* 1986), Hendrie River Water Cave (Michigan) (Volz & Yao 1991), West Virginia Caves (Rutherford & Huang 1994), and Kozlov Rob Cave (Slovenian) (Gunde-Cimerman *et al.* 1998).

Other fungal species recorded in RCCP, particularly in wet soil and ponds formed when the river rises, included various species of Oomycota (e.g., *Achlya americana* Humphrey, *A. parasitica* Coker, *Brevilegnia* sp., and *Saprolegnia* sp.) Nieves-Rivera & Betancourt-López (1994) isolated *Saprolegnia ferax* (Gruith.) Thuret and *Achlya americana* Humphrey from a stream pond in El Convento Cave Spring System (Guayanilla, southwestern Puerto Rico), an ancient course of Río Camuy (Troester 1988). A chytrid (*Rhizophydium* sp.) was isolated from Tres Pueblos Sinkhole in 1993 and also has been isolated from a stream (Quebrada de Oro) at Mayagüez (Nieves-Rivera, unpubl. data).

In the entrances of RCCS, several higher fungal species, *Cotylidia aurantica* (Pers.) A. L. Welden, *Rigidoporus microporus* (Fr.) Overeem., *Ramaria* sp., *Geastrum* cf. *saccatum* Fr. (Fig. 3J), *Marasmius* cf. *atrorubens* (Berk.) Berk., and *Polyporus tenuiculus* (Beauv.) Fr. [= *Favolus brasiliensis* (Fr.)

Fr.] were collected. *Polyporus tenuiculus* was observed being consumed by the land snail *Caracolus caracola* L. (Pulmonata: Gastropoda). Lodge (1996) reported *C. caracola* devouring basidiocarps of *P. tenuiculus* from El Verde Long Term Ecological Research site in the Luquillo Mountains, northeastern Puerto Rico. Undetermined microfungi are dominant on bat-guano deeper in RCCS (e.g., Cathedral Cave). These mycelial mats are found in caves with bat colonies in the northern and southern karst areas of Puerto Rico (Nieves-Rivera, unpubl. data).

Myxomycetes (plasmodial slime molds) *Arcyria* cf., *Comatricha* cf., and *Stemonitis* cf. *herbatica* Peck were collected from all entrances sites, except Empalme Sinkhole. Eight species of dictyostelids (cellular slime molds) were isolated from bat-guano enriched soil, or leaf litter, including *Dictyostelium aureostipes* Cavender, Raper & Norberg, *D. citrinum* Vadell, Holmes & Cavender, *D. giganteum* Singh, *D. macrocephalum* Hagiwara, Yeh & Chien, *D. mucoroides* Brefeld, *D. purpureum* Olive, *Polysphondylium pallidum* Olive, and *P. violaceum* Brefeld (Table 1). *Polysphondylium violaceum* and an undetermined myxomycete plasmodium were previously isolated from bat-guano enriched soil from El Caballo/Pájaros Cave (Playa de Pájaros) and a bat cave (Uvero), both in Mona Island (Landolt, pers. comm. 1998; Nieves-Rivera & McFarlane 2001: map). *Polysphondylium violaceum* is fairly widespread in many tropical, subtropical, and temperate habitats (Landolt *et al.* 1992; Reeves *et al.* 2000); *P. violaceum* has been isolated from Appalachian Mountains and eastern Russia, and is widely distributed at lowlands and high altitudes – one isolate came from ~2000 m

msl (Landolt, pers. comm. 2002). *Dictyostelium* and *Polysphondylium* species are inhabitants of moist forest soils and litter (Stephenson & Landolt 1996; Stephenson *et al.* 1999).

Forty-four of the 50 fungi and slime molds are new records for Cathedral Cave, Clara Cave, and Empalme and Tres Pueblos Sinkholes of RCCP (part of the RCCS). The cellular slime molds *Dictyostelium citrinum* and *D. macrocephalum* are new records for Puerto Rico.

ACKNOWLEDGMENTS

I thank John C. Landolt (Shepherd College, Shepherdstown, WV) for the isolation and identification of cellular slime molds. Further information was kindly granted by Walter A. Cardona Bonet, Juan R. Carvajal-Zamora, Gustavo A. Rodríguez, Stewart B. Peck, Carlos J. Santos-Flores, and Steven L. Stephenson. I appreciate the enthusiasm and cooperation of the staff of the National Parks Company of the Commonwealth of Puerto Rico, in particular Ángel D. López (director) and Sonia Decós (secretary). I am indebted to the guides Nancy N. Mercado and José L. Martínez for their valuable assistance. I thank Joel L. Colón, Luis E. Collazo, William Lozada, Yanaliz Lugo, Ancizar Lugo, Carlos G. Muñoz, Sylvia E. Ramos, Augusto C. Carvajal-Vélez, José De Santiago, Rosario Gaud, José A. Rubiano, and Lydia I. Rivera-Vargas for assistance with field and laboratory work. The comments and suggestions made to the text by various colleagues and reviewers are most appreciated. This project was partially supported by the University of Puerto Rico Sea Grant College Program Budget (90-4). Photographs included in Figures 2A-C were taken by José J. Vargas and Peter Rocafort digitized the figures.

REFERENCES

- Alexopoulos, C.J., Mims, C.W., & Blackwell, M., 1996, *Introductory mycology*, 4th ed.: New York, Wiley & Sons, Inc.
- Beck, B.F., Fram, M., & Carvajal-Zamora, J.R., 1976, *The Aguas Buenas Caves, Puerto Rico: geology, hydrology and ecology with special reference to the histoplasmosis fungus*: National Speleological Society Bulletin, v. 38, p. 1-16.
- Betancourt, C., Pérez, O., Seguí, D., & Emmanueli, I., 1988, *Evaluación micológica de la Cueva Ensueño, Hatillo, Puerto Rico* [Unpubl. report]: Mayagüez, Puerto Rico, Estudio Técnico para la Compañía Lebrón & Associates, 14 p.
- Boucias, D.G., Pendland, J.C., & Latge, J.P., 1988, *Nonspecific factors involved in attachment of entomopathogenic deuteromycetes to host insect cuticle*: Applied and Environmental Microbiology, v. 54, p. 1795-1805.
- Cadilla, J.F., 1962, *Informe sobre la existencia de guano animal en la parte sur de Puerto Rico* [Unpubl. report]: San Juan, Puerto Rico, Administración de Fomento Económico de Puerto Rico, Laboratorios de Investigaciones Industriales.
- Campbell, M.T., 1994, *The guano trade: Pre-World War I connection between Puerto Rico and Germany*, in 19th Caribbean Studies Association: Mérida, México, Caribbean Studies Association.
- Carvajal-Zamora, J.R., & Nieves-Rivera, Á.M., 1998, *Preliminary checklist of cave mycobiota of Puerto Rico with special reference to bat-guano enriched soil hyphomycetes*: Mycological Society of America/American Bryological and Lichenological Society 1998 Annual Meeting, San Juan, Puerto Rico, Abstracts in Inoculum, v. 49, n. 2, p. 13.
- Conde Costas, C., & González, C., 1990, *Las cuevas y cavernas en el bosque xerofítico de Guánica*: Acta Científica, v. 4, p. 113-126.
- Cubbon, B.D., 1976, *Cave flora*, in Ford, T.D., & Cullingford, C.H.D. (eds.), *The science of speleology*: London, Academic Press, p. 423-452.
- Culver, D.C., 1982, *Cave life. Evolution and ecology*: Cambridge & London, Harvard University Press.
- Del Llano, M., 1982, *La vegetación de las Cuevas del Río Camuy*, in Díaz-Díaz, C.A. (ed.), *La Cueva Clara de Empalme. Aspectos fundamentales para su desarrollo turístico* [Unpubl. report]: San Juan, Puerto Rico, Area de Investigaciones Científicas, Departamento de Recursos Naturales, p. 41-56.
- Díaz-Díaz, C.A., 1982, *La Cueva Clara de Empalme. Aspectos fundamentales para su desarrollo turístico* [Unpubl. report]: San Juan, Puerto Rico, Area de Investigaciones Científicas, Departamento de Recursos Naturales.
- Díaz-Díaz, C.A., 1983, *Los mamíferos de Puerto Rico*, in Vivaldi, J.L. (ed.), *Compendio enciclopédico de los recursos naturales, v. 1*: San Juan, Puerto Rico, Departamento de Recursos Naturales, p. 1-221.
- Ewel, J.J., & Whitmore, J.L., 1973, *The ecological life zones of Puerto Rico and the U.S. Virgin Islands*: U.S. Forest Service, Institute of Tropical Forestry Research Publication ITF-18.
- Frank, E.F., 1998, *History of the guano mining industry, Isla de Mona, Puerto Rico*: Journal of Caves and Karst Studies, v. 60, p. 121-125.
- Gaur, P.K., & Lichtwardt, R.W., 1980, *Preliminary visual screening of soil samples for the presumptive presence of Histoplasma capsulatum*: Mycologia, v. 72, p. 259-269.
- Gillieson, D., 1996, *Caves. Processes, development and management*: Cambridge, Massachusetts: Blackwell Publishers.
- Gunde-Cimerman, N., Zalar, P., & Jeram, S., 1998, *Mycoflora of cave cricket Trogliphilus neglectus cadavers*: Mycopathologia, v. 141, p. 111-114.
- Gurnee, R.H., 1967, *The Río Camuy Cave Project, Puerto Rico*: Bulletin of the National Speleological Society, v. 29, p. 27-34.
- Gurnee, R.H., & Gurnee, J., 1974, *Discovery at Río Camuy*: New York, Crown Publishers.
- Gurnee, R.H., & Gurnee, J., 1987, *Río Camuy Cave Park, Puerto Rico. Open to the public*: National Speleological Society News, v. 45, n. 5, p. 1-11.
- Hawksworth, D.L., Kirk, P.M., Sutton, B.C., & Pegler, D.N., 1995, *Ainsworth & Bisbi's Dictionary of fungi*: Cambridge, CAB International & Cambridge University Press.
- Hoff, G.L., & Bigler, W.J., 1981, *The role of the bats in the propagation and spread of histoplasmosis: A review*: Journal of the Wildlife Diseases, v. 17, p. 191-196.
- Hoffman, A., Palacios-Vargas, J.G., & Morales-Malacara, J.B., 1986, *Manual de bioespeleología (con nuevas aportaciones de Morelos y Guerrero, México)*: México, D.F., Universidad Nacional de México.
- Holsinger, J.R., & Peck, S.B., 1967, *A new genus and species of subterranean amphipod (Gammaridae) from Puerto Rico, with notes on its ecology, evolution and relationship to other Caribbean amphipods*: Crustaceana, v. 15, p. 249-262.
- Landolt, J.C., Stephenson, S. L., & Stihler, C.W., 1992, *Cellular slime molds in West Virginia caves including notes on the occurrence and distribution of Dictyostelium rosarium*: Mycologia, v. 84, p. 399-405.
- Lewis, W.C., 1989, *Histoplasmosis, a hazard to new tropical cavers*: National Speleological Society Bulletin, v. 51, p. 52-65.
- Lodge, D.J., 1996, *Microorganisms*, in Reagan, D.P., & Waide, R.B. (eds.), *The food web of a tropical rain forest*: Chicago, University of Chicago Press, p. 53-108.
- Lugo, A.E., Miranda-Castro, L., Vale, A., del Mar-López, T., Hernández-Prieto, E., García-Martinó, A., Puente-Rolón, A.R., Tossas, A.G., McFarlane, D.A., Miller, T., Rodríguez, A., Lundberg, J., Thomlinson, J., Colón, J., Schellekens, J., Ramos, O., & Helmer, E., 2001, *Puerto Rican karst— a vital resource*. USDA Forest Service General Technical Report WO-65, p. 1-100.
- Martínez-Quendo, J.A., 1983, *Sistema de Cavernas del Río Camuy: Concepto de un desarrollo para la recreación y su conservación*: Revista de Planificación ANÁLISIS, v. 2, n. 3, p. 34-35.
- Monroe, W.H., 1976, *The karst landforms of Puerto Rico*: U.S. Geological Survey Professional Paper 899, p. 1-69.

- Monroe, W.H., 1980, Geology of the middle Tertiary formations of Puerto Rico: Washington, D.C., Department of the Interior, U.S. Geological Survey.
- Nieves-Rivera, Á.M., & Betancourt-López, C., 1994, *Saprolegnia ferax* (Gruith.) Thuret (Oomycetes), new record for Puerto Rico: Caribbean Journal of Science, v. 30, p. 288-290.
- Nieves-Rivera, Á.M., & Carvajal-Zamora, J.R., 2000, Cave mycobiota of Puerto Rico, with special reference to those inhabiting bat-guano enriched soil: Memorias XIX Simposio Flora y Fauna del Caribe, Universidad de Puerto Rico, Humacao, Abstracts, p. 17.
- Nieves-Rivera, Á.M., & McFarlane, D.A., 2001, In search of the extinct hutia in cave deposits of Isla de Mona, P.R.: National Speleological Society News, v. 59, p. 92-95.
- Nieves-Rivera, Á.M., Betancourt, C., & Mignucci, J.S., 1999, Hymenomyces and gasteromyces (Basidiomycotina) of Mona Island Commonwealth Reserve, Puerto Rico: University of Puerto Rico Agricultural Experiment Station Bulletin, v. 298, p. 1-91.
- Orpurt, P.A., 1964, The microfungus flora of bat cave soils from Eleuthera Islands, The Bahamas: Canadian Journal of Botany, v. 42, p. 1629-1633.
- Peck, S.B., 1974, The invertebrate fauna of tropical American caves, part II. Puerto Rico, an ecological and zoogeographical analysis: Biotropica, v. 6, p. 14-31.
- Peck, S.B., 1981, Zoogeography of invertebrate cave faunas in southwestern Puerto Rico: National Speleological Society Bulletin, v. 43, p. 70-79.
- Peck, S.B., 1994, A synopsis of the invertebrate cave fauna of Puerto Rico, Mona Island and the Virgin Islands, in Juberthie, C., and Decu, V. (eds.), Encyclopaedia Biospeologica, v. 43: Moulis-Bucarest, p. 505-512.
- Reeves, W.K., Jensen, J.B., & Ozier, J.C., 2000, New faunal and fungal records from caves in Georgia, USA: Journal of Caves and Karst Studies, v. 62, p. 169-179.
- Reyes-Colón, C.M., 1999, Bryoflora of limestone sinkholes in the north-central karst zone of Puerto Rico [MSc thesis]: Mayagüez, Puerto Rico, University of Puerto Rico.
- Reyes-Colón, C.M., & Sastre-DJ, I., 2000, Bryophytes of two sinkholes of the north-central karst zone of Puerto Rico: Evansia, v. 17, p. 68-76.
- Rodríguez-Durán, A., 1998, Non-random aggregations and distribution of cave-dwelling bats in Puerto Rico: Journal of Mammalogy, v. 79, p. 141-146.
- Rodríguez-Durán, A., & Lewis, A.R., 1987, Patterns of population size, diet and activity time for a multispecies assemblage of bats at a cave in Puerto Rico: Caribbean Journal of Science, v. 23, p. 352-360.
- Roselló, J.G., Mari Mutt, J.A., & Betancourt, C., 1986, Listado de esporas de hongos imperfectos ingeridos por 10 especies de colémbolos colectados de hierbas en el campus del Recinto Universitario de Mayagüez: Caribbean Journal of Science, v. 22, p. 115-121.
- Rutherford, J.M., & Huang, L.M., 1994, A study of fungi of remote sediments in West Virginia caves and a comparison with reported species in the literature: National Speleological Society Bulletin, v. 56, p. 38-45.
- Santos-Flores, C.J., 2001, The taxonomy and distribution of the freshwater micro-crustaceans and "green algae" of Puerto Rico, with a synopsis on West Indian Limnology and three contributions to American Cladocerozoology [PhD dissert.]: Madison, Wisconsin, University of Wisconsin-Madison.
- Santos-Flores, C.J., & Betancourt-López, C., 1997, Aquatic and water-borne hyphomycetes (Deuteromycotina) in streams of Puerto Rico (including records from other neotropical locations): Caribbean Journal of Science Special Publication 2.
- Sparrow, Jr., F.K., 1960, Aquatic phycomycetes, 2nd ed.: Ann Arbor, Michigan, The University of Michigan Press.
- Stephenson, S.L., & Landolt, J.C., 1996, The vertical distribution of dictyostelids and myxomycetes in the soil/litter microhabitat: Nova Hedwigia, v. 62, p. 105-117.
- Stephenson, S.L., Landolt, J.C., & Moore, D.L., 1999, Protostelids, dictyostelids, and myxomycetes in the litter microhabitat of the Luquillo Experimental Forest, Puerto Rico: Mycological Research, v. 103, p. 209-214.
- Stevenson, J.A., 1975, Fungi of Puerto Rico and the American Virgin Islands: Ann Arbor, Michigan, Braun-Brumfield, Inc.
- Sullivan, B.G.N., 1966, A biologist goes underground to study animal life: Animal Kingdom [October], p. 130-139.
- Sullivan, B.G.N., 1967, Ecology of the Río Camuy Cave area, Puerto Rico: Bulletin of the National Speleological Society, v. 29, p. 38-39.
- Thraillkill, J.V., 1967, Geology of the Río Camuy Cave area, Puerto Rico: Bulletin of the National Speleological Society, v. 29, p. 35-37.
- Torres-González, A., 1983, Hydrologic study of the Río Camuy Cave System, Puerto Rico [MSc thesis]: Mayagüez, Puerto Rico, University of Puerto Rico.
- Troester, J.W., 1988, Geology of the Cueva Ensueño area, in Gurnee, J. (ed.), The Ensueño Cave study: New Jersey, National Speleological Foundation, Inc., p. 18-21.
- Troester, J.W., 1994, The geochemistry, hydrogeology, and geomorphology of the Río Camuy drainage basin, Puerto Rico, a humid tropical karst [PhD dissertation]: Pennsylvania, Penn State University.
- Troester, J.W., & White, W.B., 1986, Geochemical investigation of three tropical karst drainage basins in Puerto Rico: Ground Water, v. 24, p. 475-482.
- Vives, M.A., Estremada, W., Deliz, J.A., Hernández, L., Álvarez, A., & Reverol, J., 1985, Estudio preliminar- Visitas # 1 y 2, Plantas del área de Cuevas de Camuy. [Unpubl. report]: Quebradillas, Puerto Rico, Club Roy C. Woodbury, Escuela Superior Quebradillas.
- Volz, P.A., & Yao, J.P., 1991, Micro-fungi of the Hendrie River Water Cave, Mackinac County, Michigan: National Speleological Society Bulletin, v. 53, p. 104-106.